



DHS Quarantine Unit Admission Standing Orders:

Patient Name: _____

Allergies: _____

Date of Birth: _____ Age: _____

Advance Directives: _____

Acuity Level: 1 2 3

Vital Signs:

- Acuity 1: TID - 8am, 2pm, 8pm.
- Acuity 2: BID - 8am & 8pm
- Acuity 3: Daily 8am

Medical providers can change the frequency of vital signs as needed for any change in patient condition.

Diet:

- Regular
- Diabetic
- Renal Diet (Low Salt/Low)
- Cardiac Diet (Low Salt/Fat)
- Other: _____

Fluids:

- No Restrictions
- Fluid Restriction < 2 L/day.
- Push PO Fluids/Gatorade

Medications:

- Acetaminophen (Tylenol) 1000 mg po every 8 hours as needed for temperature > 100.4.
- Acetaminophen (Tylenol) 650 mg po every 6 hours as needed for mild-moderate pain.
- Magnesium Hydroxide (MOM) 30 ml/po every 12 hours as needed for constipation.
- Diphenhydramine HCL (Benadryl) 25/50 mg PO every 6 hours as needed for itching, sleep or allergy.
- Menthol (Halls) one lozenge PO every 2 hours as needed for sore throat.
- Calcium Carbonate (TUMS) 1500 mg PO every 6 hours for indigestion/heartburn as symptoms occur (no more than 10 tabs in a 24-hour period) **Please do not give if ESRD.**
- Mylanta 10-20 ml every 6 hours as needed for indigestion/heartburns as symptoms occur (no more than 10 tabs in a 24-hour period) **Please do not give if ESRD.**
- Guaifenesin Syrup (Tussin) 400 mg (20 ml) PO every 4 hours as needed for cough.
- Aspirin (ASA) 81 mg Pox 1 if reports of Chest Pain and call 911.
- Naloxone (Narcan) 0.4 mg/ml IM if signs & symptoms of Opiate Overdose; if no response, repeat x 1.
- Epinephrine (Epipen) 0.3 mg/SQ at signs and symptoms of Anaphylaxis.
- Triple antibiotic ointment PRN cut or abrasion (please check with provider).
- Ondansetron 4-8 mg every 8 hours as needed for nausea & vomiting.
- Imodium 2 mg PO every 8 hours for Diarrhea (unless suspect for C-Diff).
- Glucagon 1 mg/IM x 1 for blood sugar <50 with symptoms of hypoglycemia and unable to take per mouth.
- Ibuprofen (Motrin) 400 mg PO every 6 hours as needed for mild-moderate pain (1-7).
- Flu shot 24 – 48 hours prior to discharge pending patient consent.
- Albuterol inhaler 1-2 puffs every 4-6 hours PRN for SOB/Wheezing.
- Melatonin 5 mg/po QHS PRN for insomnia.

Other Orders:

1. For patients with Alcohol use disorder who decline;
Benzodiazepine/Gabapentin Taper (See MAT Protocol) and are at risk for exhibiting withdrawal symptoms:
 - ___ Beer ever ___ Hours/PRN
 - ___ Oz every ___ Hours/PRN
2. For patients with nicotine use disorder who decline nicotine replacement therapy (See MAT Protocol) and understand risks of continued smoking:
 - ___ Cigarettes every ___ hours PRN.
3. For Patient who will not stay in isolation/Quarantine without Marijuana:
 - ___ Edible every 8/12 hours PRN
 - CBD 1 capsule q 12 Hours/PRN
 - * No more than 2 cap/ 24 hours
4. Other:
 - *See MAT protocol for Opiates overdose
 - V/S per protocol.
 - Anticipate D/C planning & Date.
 - See MD medication orders.
 - _____
 - _____
 - _____

Provider Name: _____

Signature: _____

Date: _____ Time: _____